



Specimen ID: 00000000000

PATIENT, ANY

Gender:

DOB: 00/00/0000

Age(y/m/d): 000/00/00

Acct #: 00000000 **Phone:** (000) 000-000

CMBP QA TEST ACCOUNT CMBP QA DEPT RTP.

RTP

RTP NC 27215-5820

Patient Details Specimen Details

 Date collected:
 00/00/0

 Date received:
 00/00/0

 Date entered:
 00/00/0

00/00/0000 00:00 Local 00/00/0000 00:00 00/00/0000 00:00 Physician Details Ordering: Referring:

ID:

 Patient ID:
 Date reported:
 00/00/0000 00:00 ET
 NPI: 0000000000

Specimen Type: Whole Blood

Clinical Indication: Testing for familial cancer risk variant(s): BRCA2 c.9812T>C

RESULTS

POSITIVE FOR TARGETED VARIANT

No clinically significant variants were identified.

At least one variant of uncertain significance (VUS) was identified; see Additional Clinical Information for details.

GENE	VARIANT
BRCA2	No pathogenic variants were identified.

INTERPRETATION

The *BRCA2* c.9812T>C (p.Leu3271Ser) targeted variant was identified in this individual. No pathogenic variants were identified.

ADDITIONAL CLINICAL INFORMATION

Variant(s) of uncertain significance (VUS) in this individual

BRCA2 c.9812T>C (p.Leu3271Ser) (VUS - uncertain significance). Insufficient information is available to classify this variant as either benign or pathogenic.

The significance of VUS in association with relative risk for HBOC (Hereditary Breast and Ovarian Cancer Syndrome) is uncertain at the time of reporting. Evidence for variant classification is available upon request.

Electronically released by: Any Physician, PhD

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Any Physician, MD,



PATIENT: PATIENT, ANY

Specimen ID: 00000000000 **CONTROL ID: DOB:** 00/00/0000 **Patient ID:** Date Collected: 00/00/0000 00:00 Local LAB CASE ID: 000000000000000

ADDITIONAL CLINICAL INFORMATION

NCCN Guidelines

When BRCA1 and BRCA2 results are negative, additional testing may be helpful for some patients with breast, ovarian, and pancreatic cancer. Guidelines from the National Comprehensive Cancer Network® (NCCN®) recommend considering germline genetic testing for high-penetrance breast and/or ovarian cancer genes (including, but not limited to, BRCA1/2, CDH1, PALB2, PTEN, and TP53) in patients with any of the criteria in the table below. To discuss comprehensive genetic testing for breast, ovarian, and pancreatic cancer genes, an Integrated Genetics' Genetic Coordinator is available at 800-345-4363.

Breast cancer diagnosed ≤ age 45	Ovarian cancer at any age
Breast cancer diagnosed age 46-50 with ≥1 close	Breast cancer diagnosed at any age, with ≥1 close relative
relative with breast, ovarian, pancreatic, or high	with breast cancer ≤ age 50, or ovarian, pancreatic, or
grade prostate cancer	metastatic prostate cancer
Male breast cancer at any age	Pancreatic cancer at any age
Triple negative breast cancer diagnosed ≤60	Metastatic prostate cancer at any age
Breast, ovarian, or pancreatic cancer at any age;	A first or second degree relative meeting any of the criteria
and Ashkenazi Jewish ancestry	in this table

RECOMMENDATIONS

Based on information provided, this patient meets National Comprehensive Cancer Network® (NCCN®) clinical quidelines that recommend considering testing for additional genes. See Additional Clinical Information for more details. Integrated Genetics' Genetic Coordinators are available at (800) 345-GENE to discuss additional genetic testing if desired.

Genetic counseling is recommended to discuss the potential clinical and/or reproductive implications of these results, as well as recommendations for testing family members. To access Integrated Genetics Genetic Counselors please visit www.integratedgenetics.com/genetic-counseling or call (855) GC-CALLS (855-422-2557).

METHODS AND LIMITATIONS

Next-generation sequencing: Genomic regions of interest are selected using a custom capture reagent for target enrichment and sequenced via the Illumina(R) next generation sequencing platform. Regions of interest include all exons and intron/exon junctions (+/-20 nucleotides) of the BRCA1 (NM_007294.3) and BRCA2 (NM_000059.3) genes. Sequencing reads are aligned with the human genome reference GRCh37/hg19 build. Minimum mean coverage is 40X. Any segment failing minimum read depth coverage is rescued by bi-directional Sanger sequencing to complete sequence analysis. Variants, including SNVs and CNVs, are identified using a custom bioinformatics pipeline.

Reported variants: Pathogenic and likely pathogenic variants and variants of uncertain significance (VUS) are reported. Non-deletion variants are specified using the numbering and nomenclature recommended by the Human Genome Variation Society (HGVS, http://www.hgvs.org/). Benign variants are not reported. Variant classification and confirmation are consistent with ACMG standards and guidelines (Richards, PMID:25741868; Rehm, PMID:23887774). Detailed variant classification information is available upon request. A variant of uncertain significance (VUS) should not be used in clinical decision making; a VUS is classified based on inadequate or conflicting evidence regarding its pathogenicity or clinical relevance.

Electronically released by: Any Physician, PhD

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Any Physician, MD.



BRCAssure® BRCA2 Targeted Analysis

PATIENT: PATIENT, ANY

DOB: 00/00/0000

 Specimen ID: 00000000000 Date Collected: 00/00/0000 00:00 Local

METHODS AND LIMITATIONS

Limitations: Technologies used do not detect germline mosaicism and do not rule out the presence of large chromosomal aberrations, including rearrangements, gene fusions, or variants in regions or genes not included in this test, or possible inter/ intragenic interactions between variants. Variant classification and/or interpretation may change over time if more information becomes available. False positive or false negative results may occur for reasons that include: genetic variants, pseudogene interference, technical handling, blood transfusions, bone marrow transplantation, mislabeling of samples, or erroneous representation of family relationships. For heterozygous variants in the same gene the assay cannot determine whether they are on the same or a different chromosomes; to determine phase and clinical significance, rarely, parental testing may be required. Exact breakpoints of exon-level deletions/duplications are not determined. The presence of an inherited cancer syndrome due to a different genetic cause cannot be ruled out. Any interpretation should be clinically correlated with information about the patient's presentation and relevant family history.

REFERENCES

- 1. NCCN Genetic/Familial High Risk Assessment: Breast, Ovarian, and Pancreatic. Version 1.2020.
- 2. Petrucelli, et al. *BRCA1* and *BRCA2*-Associated Hereditary Breast and Ovarian Cancer. GeneReviews, updated 2016. PMID: 20301425.

This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

Electronically released by: Any Physician, PhD

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Any Physician, MD.