

Specimen ID: \_\_\_\_\_  
 Control ID: \_\_\_\_\_ Lab Case ID: \_\_\_\_\_

Acct #: \_\_\_\_\_

Phone: \_\_\_\_\_

**REPORT, SAMPLE**

Patient Details	Specimen Details	Physician Details
DOB:	Date collected:	Ordering:
Age(y/m/d):	Date received:	Referring:
Gender:	Date entered:	ID:
Patient ID:	Date reported:	NPI:

**Specimen Type:** Whole Blood

**Clinical Indication:** Testing for familial cancer risk variant(s): BRCA2 c.3785C>G

**RESULTS**

**NEGATIVE FOR TARGETED VARIANT(S)**

No clinically significant variants or variants of uncertain significance were identified.

GENE	VARIANT
BRCA2	<b>NEGATIVE</b> No pathogenic variants were identified.

**INTERPRETATION**

The BRCA2 c.3785C>G (p.Ser1262X) targeted variant was NOT identified in this individual. No pathogenic variants were identified.

**ADDITIONAL CLINICAL INFORMATION**

**NCCN Guidelines**

When BRCA1 and BRCA2 results are negative, additional testing may be helpful for some patients. Guidelines from the National Comprehensive Cancer Network® (NCCN®) recommend considering germline genetic testing for additional breast, ovarian, prostate, and/or pancreatic cancer susceptibility genes in patients meeting any of the criteria in the table below. To discuss comprehensive genetic testing, a Labcorp Genetic Coordinator is available at 800-345-4363.

Breast cancer diagnosed ≤ age 50
Ovarian cancer, pancreatic cancer, male breast cancer, or metastatic/high-risk prostate cancer at any age
Breast cancer diagnosed at any age and one of the following: <ul style="list-style-type: none"> <li>To aid in PARP inhibitor or olaparib treatment</li> <li>Triple negative breast cancer</li> <li>Multiple primary breast cancers</li> <li>≥ 3 total diagnoses of breast cancer in patient and/or close blood relatives</li> </ul>
Breast or prostate cancer diagnosed at any age and one of the following: <ul style="list-style-type: none"> <li>≥ 1 close relative with breast cancer ≤ age 50 or with triple negative breast cancer at any age</li> <li>≥ 1 close relative with ovarian, pancreatic, male breast, or metastatic/high-risk prostate cancer</li> <li>≥ 2 close relatives with breast or prostate cancer at any age</li> <li>Ashkenazi Jewish ancestry</li> </ul>
Patients with a first or second degree relative meeting certain criteria in this table may consider germline genetic testing as well. Complete criteria may be found at NCCN.org.

**Electronically released by: Max Scott**

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Anjen Chenn, M.D., Ph.D.

Date Reported: \_\_\_\_\_

**FINAL REPORT**

Page 1 of 3

This document contains private and confidential health information protected by state and federal law. If you have received this document in error, please call 800-222-7566.

©1995-2023 Laboratory Corporation of America® Holdings. All rights reserved.

Patient:

DOB:

Patient ID:

Control ID:

Lab Case ID:

Specimen ID:

Date Collected:

## RECOMMENDATIONS

Genetic counseling is recommended to discuss the potential clinical and/or reproductive implications of these results, as well as recommendations for testing family members. To access Labcorp Genetic Counselors please visit <https://womenshealth.labcorp.com/genetic-counseling> or call (855) GC-CALLS (855-422-2557).

## METHODS AND LIMITATIONS

**Next-generation sequencing:** Genomic regions of interest are selected using a custom capture reagent for target enrichment and sequenced via the Illumina(R) next generation sequencing platform. Regions of interest include all exons and intron/exon junctions (+/-20 nucleotides) of the *BRCA1* (NM\_007294.3) and *BRCA2* (NM\_000059.3) genes. Regions of interest may be extended to include intronic sequences known to harbor pathogenic/likely pathogenic variants. Sequencing reads are aligned with the human genome reference GRCh37/hg19 build. Minimum mean coverage is 40X. Any segment failing minimum read depth coverage is rescued by bi-directional Sanger sequencing to complete sequence analysis. Variants, including SNVs and CNVs, are identified using a custom bioinformatics pipeline.

**Reported variants:** Pathogenic and likely pathogenic variants and variants of uncertain significance (VUS) are reported. Non-deletion variants are specified using the numbering and nomenclature recommended by the Human Genome Variation Society (HGVS, <http://www.hgvs.org/>). Benign variants are not reported. Variant classification and confirmation are consistent with ACMG standards and guidelines (Richards, PMID:25741868; Rehm, PMID:23887774). Detailed variant classification information is available upon request. A variant of uncertain significance (VUS) should not be used in clinical decision making; a VUS is classified based on inadequate or conflicting evidence regarding its pathogenicity or clinical relevance.

**Limitations:** Technologies used do not detect germline mosaicism and do not rule out the presence of large chromosomal aberrations, including rearrangements, gene fusions, or variants in regions or genes not included in this test, or possible inter/ intragenic interactions between variants. Variant classification and/or interpretation may change over time if more information becomes available. False positive or false negative results may occur for reasons that include: genetic variants, pseudogene interference, technical handling, blood transfusions, bone marrow transplantation, mislabeling of samples, or erroneous representation of family relationships. For heterozygous variants in the same gene the assay cannot determine whether they are on the same or different chromosome; to determine phase and clinical significance, rarely, parental testing may be required. Exact breakpoints of exon-level deletions/duplications are not determined. The presence of an inherited cancer syndrome due to a different genetic cause cannot be ruled out. Any interpretation should be clinically correlated with information about the patient's presentation and relevant family history.

## REFERENCES

1. NCCN Genetic/Familial High Risk Assessment: Breast, Ovarian, and Pancreatic. Version 2.2023.
2. Petrucelli, et al. *BRCA1*- and *BRCA2*-Associated Hereditary Breast and Ovarian Cancer. GeneReviews, updated 2022. PMID: 20301425.

This test was developed and its performance characteristics determined by Labcorp. It has not been cleared or approved by the Food and Drug Administration.

### Electronically released by: Max Scott

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Anjen Chenn, M.D., Ph.D.

Date Reported:

**FINAL REPORT**

Page 2 of 3

This document contains private and confidential health information protected by state and federal law.  
If you have received this document in error, please call 800-222-7566.

©1995-2023 Laboratory Corporation of America® Holdings.  
All rights reserved.

Patient:

Control ID:

Specimen ID:

DOB:

Patient ID:

Lab Case ID:

Date Collected:

### PERFORMING LABS

Component Type	Performed At	Laboratory Director
Technical component, processing	Laboratory Corporation of America, 1912 TW Alexander Drive, RTP, NC, 27709-0150	Anjen Chenn, MD, PhD
Technical component, analysis	Laboratory Corporation of America, 1912 TW Alexander Drive, RTP, NC, 27709-0150	Anjen Chenn, MD, PhD
Professional component	Laboratory Corporation of America, 1912 TW Alexander Drive, RTP, NC, 27709-0150	Anjen Chenn, MD, PhD

For inquiries, the physician may contact the lab at 800-345-4363

SAMPLE

**Electronically released by: Max Scott**

Testing Performed at Laboratory Corporation of America Holdings, 1912 T.W. Alexander Drive, RTP, NC 27709-0150, 1-800-345-4363, Medical Director: Anjen Chenn, M.D., Ph.D.

Date Reported:

**FINAL REPORT**

Page 3 of 3

This document contains private and confidential health information protected by state and federal law.  
If you have received this document in error, please call 800-222-7566.

©1995-2023 Laboratory Corporation of America® Holdings.  
All rights reserved.